

# Rogitamine

Phentolamine mesylate 10 mg / 1 ml

## Composition:

10 mg Phentolamine mesylate in 1 ml

## Indication:

· Impotence Therapy

## Dose:

- 1 ml bimix (30 mg Papaverine + 1 mg Phentolamine)
- 1 ml trimix (30 mg Papaverine + 1 mg Phentolamine + 10 microg PGE1)
- Take 2-10 min to work & you should attempt inter course within 2hrs after injecting the medicine.
- Not more than one dose in day.
- Not more than 2 days in a row.
- Not more than 3 times a week.

## Package:

Pack Containing 5 Ampoules

## References:

1) 1998 - 2009 Health communities.com, Inc. All rights Reserved. 2) The Journal of clinical pharmacology and New Drugs, 1971; 11: 197 - 203. 3) Department & Radiology King's collage hospital, Denmark Hill, London SE5 9RS ROYAUME - UNI. 4) Bella AJ, Brock GB. Intracavernous pharmano therapy for erectile dysfunction. Endocrine. 2004 Mar - Apr; 23(2-3): 149 - 55. 5) American Vien and Aesthetics Institute © 2002 - 2009 / All rights reserved Tuesday, May 12, 2009. 6) Meinhardt W, de la fuente RB, Lycklama aNDelechi AAB, et al. Prostaglandin E2 with Phentolamine for the treatment of erectile dysfunction. Int J impot Res 1996; 8: 1-1

For further information please contact Egypharma Scientific office: 24 Abbas elakkad St., Nasr city, Cairo


Mob.: +20 01000 32 560

Tel & Fax: +202 228 71 991

E-Mail: [egypharmaco@yahoo.com](mailto:egypharmaco@yahoo.com)



## Intra cavernosal Injection of **Rogitamine**® for



Intracavernous injection with phentolamine and papaverine appears to be a safe and effective treatment of impotence<sup>(4)</sup>

The pharmacological treatment has been successful in producing an erection in over 80% of the patients in the Erectile Dysfunction unit. This erection is more than adequate to perform intercourse. The rest of the patients in whom the injection has no effect are candidates for other forms of treatment<sup>(5)</sup>

Phentolamine in combination with other agents such as PGE1 and vasoactive intestinal peptide is used with success in the treatment of impotence.<sup>(6)</sup>

Intra Cavernosal injection (ICI) therapy is the most effective non surgical treatment for E.D. and offers an effective alternative to oral therapy.

### Erectile Dysfunction:

E.D. is defined as the persistent or repeated inability to achieve and /or sustain an erection sufficient for satisfactory sexual performance.

